CONSENT FOR EXTRACTION OF TEETH AND ANESTHESIA				
Hooman	Adamous			
Patient's I	July 15, 2020			
Patients	Name Date			
	the right to be informed about your diagnosis and planned surgery so that you can decide have a procedure or not after knowing the risks and benefits.			
My diagno	osis is:			
My planne	ed treatment is:			
Alternativ	e treatment methods include:			
All surger	ries have some risks. They include the following and others:  Swelling, bruising and pain.			
2.	Stretching of the corners of the mouth that may lead to cracking or bruising.			
3.	Possible infection that might need more treatment.			
4.	Dry socket - jaw pain that begins a few days after surgery, that may need more care.			
5.	Possible damage to other teeth close to the ones being taken out, more often those with large fillings or caps.			
6.	Numbness, pain, or changed feelings in the teeth, gums, lip, chin and/or tongue (including possible loss of taste). This is due to the closeness of tooth roots (mainly with wisdom teeth) to the nerves which can be injured or damaged. Usually the numbness or pain goes away, but in some cases, it may need more treatment or may be permanent.			
7.	Trismus – Limited opening. This is most common after wisdom teeth are taken out. Sometimes it happens because of jaw joint (TMJ) problems already there. Damage can occur to the ligaments of the jaw joint (TMJ) from having your mouth open wide and/or for a period of time. This is more common if you already have symptoms or signs.			
8.	Bleeding – oozing can often happen for several hours, but a lot of bleeding is not common.			
9.	Sharp ridges or bone splinters may form later at the edge of the hole where the tooth was taken out. These may need another procedure to smooth or remove.			
10.	Sometimes tooth roots may be left in to avoid harming important things such as nerves or a sinus (a hollow place above your upper back teeth).			
11.	The roots of the upper back teeth are often close to the sinus and sometimes a piece of root can get into the sinus. An opening may occur from the sinus into the mouth that may need			

\_12. It is very rare that the jaw will break, but it is possible in cases where the teeth are **buried very deep** in their sockets.

## **FOR FEMALE PATIENTS**

more treatment.

13.	I have told my doctor that I use birth control pills. My doctor has told me that some antibiotics and other medications may reduce the preventive effect of birth control pills, and I could conceive and become pregnant. I agree to discuss with my personal doctor using other forms of birth control during my treatment, and to continue those methods until my personal doctor says that I can stop them and use only oral birth control pills.
<b>ANESTHES</b>	SIA
I have had t	he opportunity to speak with Dr. Adamous about my options for anesthesia. These options
	al Anesthesia, Nitrous Oxide/Oxygen Analgesia with Local Anesthesia, Intravenous Sedation,
or Deep Sed	dation. After this discussion, I have chosen to have IV sedation as my anesthesia. I
understand	the risks and potential complications of anesthesia to include:
14.	Discomfort, swelling or bruising where the drugs are placed into a vein.
15.	Vein irritation, called phlebitis, where the drugs are placed into a vein. Sometimes this may grow to a level of discomfort or disability where it may be difficult to move my arm or hand.
	Sometimes medication or other treatment may be needed.
16.	Nerves travel next to the blood vessels where the drugs are placed into a vein. If the needle hits a nerve or if drugs or fluid leaks out of the vessel around a nerve, I may have numbness or pain in the nerve where it runs along the arm. Usually the numbness or pain goes away, but in some cases, it may be permanent.
17.	Allergic reactions (previously unknown) to any of the medications used.
18.	Nausea and vomiting, although not common, are possible unfortunate side effects.
19.	Conscious sedation and deep sedation/general anesthesia are serious medical procedures and, whether given in a hospital or office, carry the risk of brain damage, stroke, heart attack or death.
20. In	situations where a breathing tube is used, I may have a sore throat, hoarseness or voice change.
MY OBLIG	ATIONS:
21.	Because anesthetic or sedative medications cause drowsiness that lasts for some time, I MUST be accompanied by a responsible adult to drive me to and from surgery, and stay with me for several hours until I am recovered sufficiently to care for myself. Sometimes the effects of the drugs do not wear off for 24 hours.
22.	During recovery time (normally 24 hours), I should not drive, operate complicated machinery or devices or make important decisions such as signing documents, etc.
23.	I must have a completely empty stomach. It is vital that I have NOTHING TO EAT OR DRINK for <b>six (6) hours</b> prior to my treatment. <u>TO DO OTHERWISE MAY BE LIFE-THREATENING.</u>
24.	<b>Unless instructed otherwise</b> , it is important that I take any regular medications (high blood pressure, antibiotics, etc.) or any medicines given to me by my surgeon <b>using only small sips of water</b> .
25.	By initialing #25 I'm indicating that I have read the previous 24 paragraphs.

CONSENT
I understand that my doctor can't promise that everything will be perfect. I understand that the treatment listed above and other forms of treatment or no treatment at all are choices I have. I have read and understand the above and give my consent to surgery and chosen anesthesia. I have given a complete

and truthful medical history, including all medicines, drug use, pregnancy, etc. I certify that I speak, read and write English. All of my questions have been answered before signing this form.

Sign Here	July 15, 2020
Patient's (or Legal Guardian's) Signature	Date
Hooman Adamous DMD Doctor's Signature	July 15, 2020 Date
Witness' Signature	July 15, 2020 Date